

SURAT KETERANGAN

Nomor: 250/UNUSA/Adm-LPPM/IV/2019

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 01 April 2019.

Judul : Family Factors Associated With Quality of Life in Pulmonary Tuberculosis Patient in Surabaya, Indonesia

Penulis : Dhian Satya Rachmawati, Nursalam, Arief Wibowo, Astrida Budiarti, Riza Agustin

Identitas : Indian Journal of Public Health Research & Development, Vol.9, No. 11, November 2018

No. Pemeriksaan : 2019.04.01.115

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 18%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 01 April 2019

Ketua LPPM,



Dr. Istas Pratomo, S.T., M.T.

NPP. 16081074

Paper

by Dhian

Submission date: 01-Apr-2019 10:36AM (UTC+0700)

Submission ID: 1103465386

File name: bercuriosis_Patients_in_Surabaya_Indonesia_-_Amelia_Pitaloka.pdf (936.78K)

Word count: 5056

Character count: 30663

ISSN-0976-0245 (Print) • ISSN-0976-5506 (Electronic)

Volume 9

Number 411 OctoberNovember-December2018 2017



Indian Journal of Public Health Research & Development

An International Journal



Website:

www.ijphrd.com

EXECUTIVE EDITOR

Prof Vidya Surwade

Associate Professor, Dr Baba Saheb Ambedkar Medical College & Hospital, Rohinee, Delhi

INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **Dr. Abdul Rashid Khan B. Md Jagar Din, (Associate Professor)**
Department of Public Health Medicine, Penang Medical College, Penang, Malaysia
2. **Dr. V Kumar (Consulting Physician)**
Mount View Hospital, Las Vegas, USA
3. **Basheer A. Al-Sum,**
Botany and Microbiology Deptt, College of Science, King Saud University, Riyadh, Saudi Arabia
4. **Dr. Ch Vijay Kumar (Associate Professor)**
Public Health and Community Medicine, University of Buraimi, Oman
5. **Dr. VMC Ramaswamy (Senior Lecturer)**
Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur
6. **Kartavya J. Vyas (Clinical Researcher)**
Department of Deployment Health Research, Naval Health Research Center, San Diego, CA (USA)
7. **Prof. PK Pokharel (Community Medicine)**
BP Koirala Institute of Health Sciences, Nepal

NATIONAL SCIENTIFIC COMMITTEE

1. **Dr. Anju Ade (Associate Professor)**
Kodaya Medical College, Raichur, Karnataka
2. **E. Venkata Rao (Associate Professor) Community Medicine**
Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa
3. **Dr. Amit K. Singh (Associate Professor) Community Medicine**
VCSG Govt. Medical College, Srinagar – Garhwal, Uttarakhand
4. **Dr. R. G. Viveki (Professor & Head) Community Medicine**
Belgaum Institute of Medical Sciences, Belgaum, Karnataka
5. **Santosh Kumar Mulage (Assistant Professor)**
Anatomy, Raichur Institute of Medical Sciences Raichur (RIMS), Karnataka
6. **Dr. Gouri Ku. Padhy (Associate Professor) Community and Family Medicine**
All India Institute of Medical Sciences, Raipur
7. **Ritu Goyal (Associate Professor)**
Anaesthesia, Sarswathi Institute of Medical Sciences, Panchsheel Nagar
8. **Dr. Anand Kalaskar (Associate Professor)**
Microbiology, Prathima Institute of Medical Sciences, AP
9. **Md. Amirul Hassan (Associate Professor)**
Community Medicine, Government Medical College, Ambedkar Nagar, UP
10. **Dr. N. Girish (Associate Professor) Microbiology, VIMS&RC, Bangalore**
11. **Dr. BR Hungund (Associate Professor) Pathology, JNMC, Belgaum**
12. **Dr. Sartaj Ahmad (Assistant Professor)**
Medical Sociology, Department of Community Medicine, Swami Vivekananda Subharti University, Meerut, Uttar Pradesh, India
13. **Dr Sumeeta Soni (Associate Professor)**
Microbiology Department, B.J. Medical College, Ahmedabad, Gujarat, India

NATIONAL EDITORIAL ADVISORY BOARD

1. **Prof. Sushanta Kumar Mishra (Community Medicine)**
GSL Medical College – Rajahmundry, Karnataka
2. **Prof. D.K. Srivastava (Medical Biochemistry)**
Jamia Hamdard Medical College, New Delhi
3. **Prof. M Sriharibabu (General Medicine)** GSL Medical College, Rajahmundry, Andhra Pradesh
4. **Prof. Pankaj Datta (Principal & Prosthodontist)**
Indraprastha Dental College, Ghaziabad

NATIONAL EDITORIAL ADVISORY BOARD

5. **Prof. Samarendra Mahapatro (Pediatrician)**
Hi-Tech Medical College, Bhubaneswar, Orissa
6. **Dr. Abhiruchi Galhotra (Additional Professor) Community and Family Medicine**, All India Institute of Medical Sciences, Raipur
7. **Prof. Deepti Pruthvi (Pathologist)** SS Institute of Medical Sciences & Research Center, Davangere, Karnataka
8. **Prof. G S Meena (Director Professor)**
Maulana Azad Medical College, New Delhi
9. **Prof. Pradeep Khanna (Community Medicine)**
Post Graduate Institute of Medical Sciences, Rohtak, Haryana
10. **Dr. Sunil Mehra (Paediatrician & Executive Director)**
MAMTA Health Institute of Mother & Child, New Delhi
11. **Dr. Shailendra Handu, Associate Professor, Pharma, DM (Pharma, PGI Chandigarh)**
12. **Dr. A.C. Dhariwal: Directorate of National Vector Borne Disease Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of India, Delhi**

Print-ISSN: 0976-0245 **Electronic-ISSN:** 0976-5506 **Frequency:** Quarterly
Four issues per volume

Indian Journal of Public Health Research & Development is a double blind peer reviewed international journal. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, and Public Health Laws and covers all medical specialties concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent and South East Asia.

The journal has been assigned International Standards Serial Number (ISSN) and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases. The journal is covered by EBSCO (USA), Embase, EMCare & Scopus database. The journal is now part of DST, CSIR, and UGC consortia.

Website : www.ijphrd.com

©All right reserved. The views and opinions expressed are of the authors and not of the Indian Journal of Public Health Research & Development. The journal does not guarantee directly or indirectly the quality or efficacy of any product or service featured in the advertisement in the journal, which are purely commercial.

2
Editor

Dr. R.K. Sharma
Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)

Printed, published and owned by

2
Dr. R.K. Sharma
Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)

Published at

Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)



Indian Journal of Public Health Research & Development

www.ijphrd.com

Contents

Volume 9, Number 11

November 2018

1. The Spread Area of Malaria Vector in Timor Island, East Nusa Tenggara Province 1
Muhammiad Kazwaini, Chatarina U, Wahyuni, Monika Noshirma
2. Developing Community Resilience as a Supporting System in the Care of People with Mental Health Problems in Indonesia 6
Retno Lestari, Ah Yusuf
3. Effects of Knowledge of Vitamin D on Attitudes toward Sun Exposure among Middle-Aged and Elderly Indonesian Adults 11
Rivan Virlando Suryadinata, Bambang Wirjatmadi, Merryana Adriani, Sri Sumarmi
4. The Effects of Age And Body Mass Index on Blood Glucose, Blood Cholesterol, and Blood Pressure in Adult Women 16
Riza Fikriana, Shrimarti Rukmini Devy
5. The Antioxidant Activity and Organoleptic Properties of Soursoop Leaf Tea (*Annona Muricata* L.) and Moringa Leaf (*Moringa Olifera* L.) in Combination with Guava Leaf (*Psidium Guajava*) 22
Roy J. Irawan, Trias Mahmudiono
6. Parent Communication Regarding Sexual and Reproductive Health of Adolescent: A Qualitative Systematic Review 27
Fransiska Imavike Fevriasanty, Oedojo Soedirham
7. Five Types of Personality and the Locus of Internal Control in relation to Preeclampsia Pregnancy 33
Lusiana Meinawati, Kusnanto Kusnanto, Oedojo Soedirham
8. The Effect of Diabetes Self-Management Education, Based on The Health Belief Model, on the Psychosocial Outcome of Type 2 Diabetic Patients in Indonesia 37
Rondhianto, Kusnanto, Soenarnatalina Melantani
9. How does the Dayak Ngaju Community Treat Malaria? A Qualitative Study on the Use of Traditional Medicine in Central Kalimantan Province, Indonesia 43
Trilianty Lestari, Soedjajadi Kemian
10. The Effect of Young Coconut Water against Morning Sickness among Women in the First Trimester of Pregnancy 48
Tri Ratna Ariestini, Whindu Purnomo
11. Factors Affecting the Incidence of Chronic Energy Deficiency among Pregnant Women Attending the Pulubala Community Health Centre 53
Zuriati Mohammad, Sri Sumarmi
12. Implementation of Birth Preparedness and Complication Readiness (BPCR) in High Risk Pregnancies 58
Ika Mardiyanti, Nursalam, Arief Wibowo

13. Access to Healthcare Facilities in Poor and Underdeveloped Areas in Nusa Tenggara Timur Province	64
<i>Yendris Krisno Syamruth, H. Kuntoro</i>	
14. The Effect of Doctor Professionalism on the Quality of Medical Services at a First Level Health Facility	70
<i>Febri Endra B. S., Stefanus Supriyanto, Rubayat Indradi, Aditya Rizka R</i>	
15. Self-Help Group Therapy: The Enhancement of Self-Care Ability and Quality of Life Among the Elderly in Bali, Indonesia	76
<i>I Wayan Suardana, Ah Yusuf, Windhi Purnomo</i>	
16. Healthy Nurses for a Quality Health Care Service: A Literature Review	81
<i>Dodi Wijaya, Nyoman Anita Damayanti</i>	
17. The Support of the Family toward Children with Autism Spectrum Disorder	86
<i>Siti Maemonah, Hamidah, Nyoman Anita Damayanti, Emung Mardiyana Hidayat, Aida Novitasari, Qorry Aina, Wina Tryas Fatima</i>	
18. Family Factors Associated with Quality of Life in Pulmonary Tuberculosis Patients in Surabaya, Indonesia	91
<i>Dhian Satya Rachmawati, Nursalam, Arief Wibowo, Astrida Budiarti, Riza Agustin</i>	
19. Parenting Style Based on the Mother's Personal Mastery and the Mother-Child Attachment in Relation to Child Feeding Disorders: A Qualitative Study	96
<i>Ni Putu Sudewi, Oedojo Soedirham, Ahmad Suryawan</i>	
20. Factors Associated with Onset to Hospital Delay among Stroke Patients in the Emergency Department	101
<i>Abdulloh Machin, Muhammad Hamdan</i>	
21. Analysis of The Influence of Hersey-Blanchard Leadership and Nurse Maturity on Caring Behaviour Performance Based on Patient Perception	106
<i>Minarni Wartiningsih, Stefanus Supriyanto, Sri Widati, Danoe Soesanto</i>	
22. The Relationship of Socio-Economic and Genetic Factors with Toddler Stunting at Kenjeran Public Health Center Surabaya	111
<i>Dwi Ernawati, Puji Hastuti, Dhian Satya Rachmawati, Ari Susanti, Christina Yuliasuti, Merina Widyastuti, Mieke Izzatul Mahmudah</i>	
23. Cultural Religiosity as the Determinant Factor of a Successful Healthy City in South Kalimantan, Indonesia	116
<i>Herawati, Shrimarti R. Devy</i>	
24. The Relationship between Response Time and Patient Survival with Emergency Treatment by the Code Blue Team	122
<i>Al Afik</i>	
25. The influence of Nurse's Knowledge Level on Behaviour Changes, Attitude and 5 Moments of Hand Hygiene Compliance	127
<i>Danoe Soesanto</i>	
26. The Role of Posyandu Cadres in Improving the Growth and Development of Toddlers in RW VII Puskesmas Mojo, Surabaya	132
<i>Emung Mardiyana Hidayat, Rini Ambarwati Indriatje, Indriatje</i>	

27. Analysis of the Implementation of Pregnancy-related Health Care Services Through the Continuum of Care Approach in Puskesmas Bukittinggi City	137
<i>Evi Hasnita, Armita Sri Azhari</i>	
28. Feeding Care Patterns of Mothers Working as Shellfish Peelers on Children's Nutritional Status at Integrated Health Posts in Coastal Areas.....	142
<i>Meiana Harfika, Zhakiyah Saraswati, Dya Sustrami, Lela Nurlela</i>	
29. The Relationship between Socioeconomic Status and Personality Type with Depression in Adolescents	147
<i>Oktavianis, Rahmi Sari Kasoema</i>	
30. Consumption Patterns, Energy Adequacy, and The Nutritional Status of Softball Players	153
<i>Ratna Candra Dewi, Bambang Wirjatmadi</i>	
31. Compliance with Smoke-Free Legislation and Associated Factors: A Serial Survey in Bali, Indonesia	159
<i>Ketut Suarjana, Artawan Eka Putra, Putu Ayu Swandewi Astuti, Ketut Hari Mulyawati, Djazuly Chalidyanto</i>	
32. The association of Pre-Pregnancy Body Mass Index (BMI) and Increased Maternal Weight in the Third Trimester of Pregnancy with Foetal Weight Estimation.....	165
<i>Siti Maimunah, Andina Firdaus Supriyanto</i>	
33. The Effectiveness of Bay Leaf Extract (<i>Syzygium Polyanthum</i>) in Inhibiting the Growth of <i>Candida Albicans</i>	170
<i>Suratih, Dewa Ayu Ketut Surinati, Dewa Made Ruspawan</i>	
34. The Presenting Symptoms as a Predictor of the Hospital Arrival Time Intervals of Patients with Acute Coronary Syndrome.....	175
<i>Tony Suharsono, Shyniatry Ayu Andhika, Ahmad Hasyim Wibisono, Tina Handayani</i>	
35. Factors Influencing the Husband's Participation in Pregnancy Care in Surabaya City, Indonesia	180
<i>Nurul Fitriyah, Windhi Purnomo, Noviasari Reksohadi</i>	
36. Exploration of the Daughters' Feelings Related to Accepting their Mother's Condition of Having Breast Cancer in East Java, Indonesia.....	185
<i>Sirli Mardianna Trishinta, Retty Ratnawati, Septi Dewi Rachmawati</i>	
37. Mapping Customers: A Case Study of a University Hospital in Indonesia	190
<i>Purwaningsih, Nyoman Anita Damayanti, Nasronudin, Thini Nurul Rochmah</i>	
38. Structural Model of the Factors Related to the Family Resilience of Stroke Patients in Indonesia	194
<i>Nikmatul Fadilah, Kusnanto, Nursalam, Minarti, Asnani</i>	
39. A Gender Analysis of Traditional Contraceptive Use in Sikka District, East Nusa Tenggara, Indonesia	200
<i>Rut Rosina Rivu, Sarci Magdalena Toy, Daniela L. A. Boeky, Conrad L. H. Folamiuk</i>	
40. The Relationship between Sexual Behavior and the Prevalence of HIV/AIDS among Homosexual Men in Bukittinggi City, Indonesia	206
<i>Nurhayati, Wahyu Salendri</i>	
41. The Nutritional Status of Children Aged 1-3 Years Old Based on Food Processing Techniques in Surabaya	211
<i>Qori'ila Saidah, Yudi Handoko, Nur Chabibah, Sri Anik Rustini, Nuh Huda, Dwi Priyanti, Dini Mei Widayanti</i>	
42. Mammary Gland in Supporting the Lactation Process: A Review	215
<i>Thontowi Djauhari Nur Subchi, Merryana Andriani</i>	

Family Factors Associated with Quality of Life in Pulmonary Tuberculosis Patients in Surabaya, Indonesia

Dhian Satya Rachmawati¹, Nursalam², Arief Wibowo³, Astrida Budiarti⁴, Riza Agustin⁴

¹Doctoral Program of Public Health, Faculty of Public Health, ²Faculty of Nursing, ³Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia; ⁴STIKes Hing Tuah Surabaya, Indonesia

Abstract

Introduction: Pulmonary TB is an infectious disease caused by *Mycobacterium Tuberculosis*. The physical and psychological changes in patients with pulmonary TB can affect the patients's quality of life. The family is a key factor in relation to supporting successful treatment and recovery. This study aims to analyse the factors related to pulmonary TB patients quality of life.

Method: This research used an analytic observational design with a cross-sectional approach. A total of 73 respondents were recruited via the family members of patients with pulmonary TB using a simple random sampling technique. The data was collected using several questionnaires on the sociodemographic characteristics, family development stage, family stress & coping, environmental data and WHOQOL-BREF. The data analysis was performed using a Chi Square test.

Result: The results of this study showed that family factors significantly influence the quality of life of patients with Pulmonary TB, including the type of family ($p=0.000$), their level of education ($p=0.000$), employment ($p=0.001$) and monthly income ($p=0.002$). Other factors including the level of stress and coping and environmental health (healthy housing) were also significantly associated with quality of life ($p<0.01$). Only the family development stage had no significant relationship with quality of life.

Conclusion: The significant family factors influence the quality of life among pulmonary TB patients, which reflects the need to strengthen the role of the family in promoting successful treatment. A family with social support from the community would help to improve the quality of life of pulmonary TB patients.

Keywords: family, pulmonary TB patients, quality of life, Indonesia

Introduction

Tuberculosis (TB) is a major communicable disease that claims 100,000 lives worldwide annually.¹ In 2017, Indonesia was ranked the third among listed countries with the highest TB burden.¹ The TB prevalence in Indonesia was estimated to be 600,000 cases with approximately 430,000 new cases per year.² The East Java Province was the top second highest district regarding the number of TB cases with a total number

of 21,606 cases in 2016.² Pulmonary TB is an infectious disease caused by rod-shaped bacteria (basil) known as *Mycobacterium Tuberculosis*.¹ Poor immune system, malnutrition, and HIV positive are among the risk factors of pulmonary TB.² According to the Indonesia Ministry of Health (MoH), the highest prevalence of pulmonary TB is among people older than 45 years old, who have a low level of education, and who are unemployed.³

TB easily infects other individuals through direct contact, coughing, sneezing, and sputum (droplet nuclei) from TB patients. As a result of living closely to TB patients, their families run the risk of TB infection. Because of worries from getting infected, the other family members may limit their contact with the TB patient which results in the individual feeling isolated, depressed and neglected.⁴ TB patients are often socially stigmatised which may affect their adherence to

Corresponding Author:

Dhian Satya Rachmawati

Doctoral Program of Public Health,

Faculty of Public Health, Universitas Airlangga,
Surabaya, Indonesia

Email: dhian.satya.rachmawati-2017@fkm.unair.ac.id

effective treatment.⁴ This psychological problem would increase the patient's stress, which affect their quality of life. The physiological changes experienced by TB patients affects their physical abilities and deteriorates their quality of life.¹ The World Health Organisation Quality of Life (WHOQOL) defines quality of life as an individual's perception of life in society in the context of the existing culture and value system, related to their goals, expectations, standards, and concerns.⁵ Quality of life is a very broad concept that is influenced by the physical condition of the individual, their psychological state, level of independence, social relations and environmental condition.⁵

Family factors are significant in relation to the TB cases. A previous study reported the significant association between TB and family-related factors, such as the number of adults in the household, having a single marital status, having a family history of TB, and living in a rented house.⁶ Understanding the family factors affecting the TB cases would improve the support and interventions required in order to promote the successful treatment of pulmonary TB. Therefore, this study aimed to investigate the relationship between family-related factors and quality of life among pulmonary TB patients.

MATERIAL AND METHOD

We employed an analytical observational design with a cross-sectional approach to investigate the relationship between family factors and the quality of life of patients with pulmonary tuberculosis in Surabaya city. This was a preliminary study carried out from March to July 2017. The study population was made up of the families of TB patients who accompanied the patient to the Perak Timur community health centre, Surabaya city, during the study period. The sampling technique used in this research was non-probability sampling using a simple random sampling technique. A total of 89 people attending the community health centre had a TB positive test. 73 people who had a family member with pulmonary TB were selected and agreed to participate in the study.

The questionnaire regarding sociodemographics, family development stage, family stress and coping, environmental data, and the WHOQOL-BREF instrument was used for the data collection. A bivariate analysis using a chi-square test was applied to test the relationship between the independent variables and quality of life.

RESULTS

The sociodemographic characteristics of the respondents have been presented in Table 1. The majority of the respondents were from a traditional family (86.3%). A traditional family, in this study, represents a nuclear family consisting of two parents and their children, while a non-traditional family reflects a single parent family or extended family.⁷ High school was the highest level of education attained by most of the respondents (39.7%). Most of the respondents worked as labourers or factory workers (65.8%), with a monthly income below IDR 1,000,000.

Table 1: Sociodemographic characteristics of the study participants, N = 73

Variables	N (%)
Type of family	
Traditional	63 (86.3)
Non traditional	10 (13.7)
Level of Education	
No schooling	6 (8.2)
Elementary	15 (20.5)
Junior school	14 (19.2)
High school	29 (39.7)
Higher education	9 (12.3)
Employment	
Factory workers	48 (65.8)
Self employed	10 (13.7)
Others	15 (20.5)
Monthly income (IDR)	
< 1,000,000	38 (52.1)
2,000,000-3,000,000	29 (39.7)
> 3,000,000	6 (8.2)

According to Duvall and Miller (1985), family life consists of eight stages, namely 1) new couple, 2) first child birth family, 3) family with pre-school children, 4) family with school children, 5) family with teenage children, 6) family with adult children, 7) middle age family, and 8) elderly family.⁸ As displayed in Table 2, most of the respondents were in the fourth family development stage (19.2%) and the fifth stage (17.8%) respectively. From their answers to the questions related to the level of stress, most of respondents had only a mild level of stress (69.9%), and none of the respondents indicated themselves as having a severe level of stress. The majority of the respondents lived in unhealthy houses (65.8%), but reported having a good quality of life (71.2%).

Table 2: Characteristics of the Respondents
Based on the Family Development Stage, Level of Stress, Environmental Health and Quality of Life

Variable	N (%)
Family development stage	
Stage 1	6 (8.2)
Stage 2	3 (4.1)
Stage 3	5 (6.8)
Stage 4	14 (19.2)
Stage 5	13 (17.8)
Stage 6	12 (16.4)
Stage 7	10 (13.7)
Stage 8	10 (13.7)
Level of stress	
Mild	51 (69.9)
Moderate	22 (30.1)
High	0 (0.0)
Environmental health	
Healthy house	25 (34.2)
Unhealthy house	48 (65.8)
Quality of Life	
Poor	21 (28.8)
Good	52 (71.2)

We assessed the family-related variables including the type of family, level of education, employment, monthly income, family development stage, level of stress, and environmental health in relation to quality of life using the chi-square test. From the analysis results shown in Table 3, we found that the type of family, level of education, and monthly income were significantly associated with quality of life. No association was found between the family development stage and quality of life ($p = 0.328$). Both the level of stress and the environmental health variables showed a significant association with the quality of life of TB patients.

Table 3: Bivariate analysis of the sociodemographic characteristics, family development stage, level of stress, environmental health, and quality of life

Variables	Quality of Life		p-value
	Poor (n; %)	Good (n; %)	
Type of family			

Traditional	12 (19.0)	51 (81.0)	0.000
Non traditional	9 (90.0)	1 (10.0)	
Level of Education			
No schooling	6 (100.0)	0	0.000
Elementary	15 (100.0)	0	
Junior school	0	4 (100.0)	
High school	0	29 (100.0)	
Higher education	0	9 (100.0)	
Employment			
Factory workers	10 (20.8)	38 (79.2)	0.001
Self employed	8 (80.0)	2 (20.0)	
Others	3 (20.0)	12 (80.0)	
Monthly income (IDR)			
< 1,000,000	5 (13.2)	33 (86.8)	0.002
2,000,000-3,000,000	15 (51.7)	14 (48.3)	
> 3,000,000	1 (16.7)	5 (83.3)	
Family development stage			
Stage 1	0 (0.0)	6 (100.0)	0.328
Stage 2	2 (66.7)	1 (33.3)	
Stage 3	1 (20.0)	4 (80.0)	
Stage 4	2 (14.3)	12 (85.7)	
Stage 5	5 (38.5)	8 (61.5)	
Stage 6	3 (25.0)	9 (75.0)	
Stage 7	4 (40.0)	6 (60.0)	
Stage 8	4 (40.0)	6 (60.0)	
Level of stress			
Mild	0	51 (100.0)	0.000
Moderate	21 (95.5)	1 (4.5)	
High	0	0	
Environmental health			
Healthy house	1 (4.0)	24 (96.0)	0.001
Unhealthy house	20 (41.7)	28 (58.3)	

Discussion

In our study, several factors including the type of family, level of education, employment, monthly income, level of stress, and environmental health were significantly associated with the quality of life of pulmonary TB patients. Chronic disease affected the physical and mental health, which in turn decreased their quality of life.⁹ The quality of life decreased along with

the emergence of the general symptoms of pulmonary TB such as coughing, fever with the exertion of sputum and mucus, and weight loss but improved after the first month of treatment.² Therefore, the family has an important role to the patients' adherence to the whole TB treatment.

The type of family is significant in relation to the quality of life among patients with TB. A traditional family is founded based on the union of parents with or without children. Support from a spouse or children may increase the motivation of pulmonary TB patients to comply the treatment. The spouse can monitor the patients in taking their medication correctly and accompany them to the health facility in order to get their disease checked. In line with our study results, a previous study found that being single parent increased the odds of TB cases by 63% compared to being in a married family.⁶ The risk of TB cases also increased along with the increased number of adults in the household. Having more than 10 adults in the household increased the risk of TB cases by 2.67%.⁶

Level of education was also a significant factor related to the quality of life among TB patients. This result supported the findings of previous studies stating that level of education had a significant relationship with quality of life among patients with TB.^{10,11} Employment is significantly associated with quality of life among the TB patients in our study. Similarly, a previous study in India reported there to be a significant relationship between employment and the quality of life scores of active TB patients after a year of successful treatment.¹² Another factor of income level was significantly related to the quality of life level. In the same vein, a previous study also reported that TB patients considered the level of income as being an important factor in improving quality of life.¹³ These two factors are relevant, as having employment would enable the family of the TB patients to receive a certain level income as a form of sustainable financial support. Having a low income constrains some families in being able to afford enough food for the whole family. Moreover, a low income household only has a limited number of choices when fulfilling the nutritional needs of the family. Lack of nutrition affects the immune system, which increases the risk of having an infectious disease.¹⁴

According to Antonovsky (1979), stress is a response or a mental state from experiencing tension caused by a stressor or unresolved circumstances.¹⁵

Having social support from the social environment would prevent the TB patients from feeling isolated and lower their level of stress.¹⁵ Quality of life has increased in line with the decreasing of the physical symptoms of TB patients. Stress from a chronic disease is not only experienced by patients with active TB, but also patients with latent TB.¹⁶ In latent TB, the TB symptoms are not visible, so the patients tend to be stressed and anxious about the given diagnosis.¹⁶ The results of a previous study in Indonesia also showed a significant relationship between social support and quality of life as reflected in the decreasing of life satisfaction felt not during the initial diagnosis, but after when undergoing the intensive phase of treatment.¹⁷

A healthy house represents good environmental health. A healthy house has an integrated physical, chemical, biological condition that enables the residents to attain optimal health.¹⁸ Therefore, a healthy house should meet requirements such as to fulfil physiological and psychological needs, and to prevent the transmission of diseases as well as accidents.¹⁸ A healthy house should have adequate lighting, either from natural or artificial light, adequate ventilation for fresh air circulation, and should enable the family members to feel comfort. It also should have a disposal system for garbage and household waste, as well as safe water and food that prevents disease transmission. Having a TB patient in the house increases the risk of TB transmission to other residents.

Therefore, having a healthy house would minimise this risk. As reported by a previous study, poor housing conditions significantly reduce the level of quality of life among patients with TB and their families.¹

Conclusion

Our study results yielded a significant relationship between all of the family related factors and the quality of life among pulmonary TB patients. The type of family significantly affects quality of life. Being educated and engaged in paid employment was significantly associated with quality of life, as was the family level of stress and coping. However the family development stage showed no association with quality of life. Poor housing condition also affects the quality of life of patients with pulmonary TB. Considering the significant family factors, several strategies to improve the quality of life among these patients and their families should be implemented.

First, the patients should have an adequate level of knowledge and understanding about the disease and

the transmission of TB bacteria, so then they follow the treatment procedures and prevent further transmission. Second, the family should continuously provide a supportive environment with the community to help the patients recover from and fight TB transmission. This effort will minimise the stigma felt by the patients and increase their self-confidence. Community social support is very important in speeding up the healing process, and will increase the dignity of the patients and their families living within the community.

Ethical Clearance: Ethical approval was granted by the School of Public Health in Airlangga University, Surabaya.

Source of Funding: Self funding.

Conflict of Interest: Nil.

References

1. World Health Organisation. *Global tuberculosis report 2016*. Geneva: World Health Organisation; 2016.
2. MoH. *Tuberculosis: temukan, obati sampai sembuh (Tuberculosis: find it, treat it)*. Jakarta: Center for Data and Information; 2016.
3. Balitbangkes-MoH. *Riset Kesehatan Dasar Tahun 2013 [The 2013 Basic Health Research]*. Jakarta: Balitbangkes MoH; 2013.
4. Courtwright A, Turner AN. Tuberculosis and Stigmatization: Pathways and Interventions. *Public Health Reports*. 2010;125(4_suppl):34-42.
5. World Health Organisation. *Whoqol-old Manual*. Copenhagen: World Health Organisation. 2006.
6. Lienhardt C, Fielding K, Sillah J, Bah B, Gustafson P, Warndorff D, et al. Investigation of the risk factors for tuberculosis: a case-control study in three countries in West Africa. *International journal of epidemiology*. 2005;34(4):914-23.
7. Andarmoyo S. *Keperawatan Keluarga Konsep Teori, Proses dan Praktik Keperawatan (Family nursing concept, theory, and the nursing practice)*. Yogyakarta: Graha Ilmu. 2012.
8. Friedmann M. *Family Nursing: Theory and Practice*. Norwalk: Appleton&Lange; 1999.
9. Ahmad N, Javaid A, Sulaiman SAS, Basit A, Afridi AK, Jaber AAS, et al. Effects of multidrug resistant tuberculosis treatment on patients' health related quality of life: Results from a follow up study. *PloS one*. 2016;11(7):e0159560.
10. Duyan V, Kurt B, Aktas Z, Duyan G, Kulkul D. Relationship between quality of life and characteristics of patients hospitalised with tuberculosis. *The International Journal of Tuberculosis and Lung Disease*. 2005;9(12):1361-6.
11. Cully JA, Graham DP, Stanley MA, Ferguson CJ, Sharafkhan A, Soucek J, et al. Quality of life in patients with chronic obstructive pulmonary disease and comorbid anxiety or depression. *Psychosomatics*. 2006;47(4):312-9.
12. Muniyandi M, Rajeswari R, Balasubramanian R, Nirupa C, Gopi P, Jaggarajamma K, et al. Evaluation of post-treatment health-related quality of life (HRQoL) among tuberculosis patients. *The International Journal of Tuberculosis and Lung Disease*. 2007;11(8):887-92.
13. Marra CA, Marra F, Cox VC, Palepu A, Fitzgerald JM. Factors influencing quality of life in patients with active tuberculosis. *Health and quality of life outcomes*. 2004;2(1):58.
14. Katona P, Katona-Apte J. The Interaction between Nutrition and Infection. *Clinical Infectious Diseases*. 2008;46(10):1582-8.
15. Hansel NN, Wu AW, Chang B, Diette GB. Quality of life in tuberculosis: patient and provider perspectives. *Quality of life research*. 2004;13(3):639-52.
16. Peddireddy V. Quality of Life, Psychological Interventions and Treatment Outcome in Tuberculosis Patients: The Indian Scenario. *Frontiers in psychology*. 2016;7:1664.
17. Pratiwi GD, Maulana Z. Hubungan Dukungan Sosial Keluarga Dengan Mekanisme Koping Pada Pasien TB Paru Yang Sedang Menjalani Proses Pengobatan Di Puskesmas Legok. STIKes Widya Dharma Husada Tangerang. 2014.
18. Prüss-Ustün A, Corvalán C. Preventing disease through healthy environments. Towards an estimate of the environmental burden of disease. Geneva: World Health Organisation. 2006.

Paper

ORIGINALITY REPORT

18%

SIMILARITY INDEX

17%

INTERNET SOURCES

3%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1

i-scholar.in

Internet Source

5%

2

[Submitted to Western Governors University](#)

Student Paper

2%

3

es.scribd.com

Internet Source

2%

4

vdocuments.site

Internet Source

1%

5

docplayer.net

Internet Source

1%

6

www.ijphrd.com

Internet Source

1%

7

[Submitted to Universitas Islam Indonesia](#)

Student Paper

1%

8

wrap.warwick.ac.uk

Internet Source

1%

9

[Submitted to London School of Hygiene and Tropical Medicine](#)

1%

10	academic.oup.com Internet Source	1%
11	cloud.stikes-senior.ac.id Internet Source	1%
12	www.wpro.who.int Internet Source	1%
13	webpac2.rot13.org Internet Source	1%
14	Submitted to Universitas Muhammadiyah Yogyakarta Student Paper	1%
15	Submitted to Universitas Jember Student Paper	1%
16	www.afro.who.int Internet Source	1%

Exclude quotes On
Exclude bibliography Off

Exclude matches < 1%