



**STIKES
HANG TUAH
SURABAYA**

THE PROCEEDING BOOK
The 2nd International Nursing Conference
STIKES HANG TUAH SURABAYA

*"Developing Cross–Cultural Understanding
and Behavior in Nursing Care Services"*

**The Proceeding Book 2nd International Nursing Conference
Developing Cross-Cultural Understanding and Behavior in Nursing Care Services**

STIKES Hang Tuah Surabaya



Hak Cipta © 2016,
Sekolah Tinggi Ilmu Kesehatan Hang Tuah Surabaya
Jl. Gadung No. 1 Surabaya
Telp./Fax: (031) 8411721
Website : <http://stikeshangtuah-sby.ac.id>
Email : info@stikeshangtuah-sby.ac.id

Hak cipta dilindungi undang-undang. Dilarang memperbanyak sebagian atau seluruh isi buku ini dalam bentuk apa pun, baik secara elektronik maupun mekanik, termasuk memfotokopi, merekam, atau dengan menggunakan sistem penyimpanan lainnya, tanpa izin tertulis dari Penerbit.

UNDANG-UNDANG NOMOR 19 TAHUN 2002 TENTANG HAK CIPTA

1. Barang siapa dengan sengaja dan tanpa hak mengumumkan atau memperbanyak suatu ciptaan atau memberi izin untuk itu, dipidana dengan pidana penjara paling lama 7 (**tujuh**) tahun dan/atau denda paling banyak **Rp 5.000.000.000,00 (lima miliar rupiah)**.
2. Barang siapa dengan sengaja menyiarkan, memamerkan, mengedarkan, atau menjual kepada umum suatu ciptaan atau barang hasil pelanggaran Hak Cipta atau Hak Terkait sebagaimana dimaksud pada ayat (1), dipidana dengan pidana penjara paling lama 5 (**lima**) tahun dan/atau denda paling banyak **Rp 500.000.000,00 (lima ratus juta rupiah)**.

Stikes Hang Tuah Surabaya

The Proceeding of 2nd International Nursing Conference
Developing Cross-Cultural Understanding and Behavior in Nursing Care Services

506 Hlm, 17 x 24 cm

ISBN : 978-602-72856-1-3

CONTENT

Greeting From Stering Commite	iv
Opening Speech Head Stikes Hang Tuah Surabaya.....	v
Steering Commite	vii
Scientific Paper Reviewer	ix
Conference Schedule	x
Plenary Speaker	
Speaker 1: Prof. Dr. Chang Ching 'Ihon	xi
Speaker 2: Dr. Lesley Dornan	xvii
Speaker 3: Dr. Katrina Breden	xxv
Speaker 4: Dr. Janny Prihastuty, S.Kep., Ns., MARS.....	
List of Oral Presentation	xliii
List of Poster Presentation	xlvi
List of Participant	xlvii
Plenary Discussion Room	

SCIENTIFIC PAPER REVIEWER

Prof. Dr. Chang Ching Thon
Universitas Malaysia, Malaysia

Dr. Lesley Dornan
Resercher, Chiang Mai University

Dr. Katrina Breden
Flinders University, Australia

Dr. Ah. Yusuf, S.Kp., M.Kes
Universitas Airlangga (UNAIR), Surabaya

Dr. Rachmat Hargono, dr., MS., M.PH
Unversitas Airlangga, Surabaya

Dr. Bambang Widjanarko Otok, M.si
Institute Teknologi Sepuluh November (ITS), Surabaya

CONFERENCE SCHEDULE

TIME SCHEDULE INTERNATIONAL CONFERENCE

Saturday, November 2016

TIME	ACTIVITY	VENUE
07.00 – 08.00	Registration Open	Ballroom Santika Hotel
08.00 – 08.15	Traditional Dance Welcome Greetings Indonesian National Anthem	Ballroom Santika Hotel
08.15 – 08.30	Opening : <ul style="list-style-type: none">• Speech from Head of STIKES Hang Tuah Surabaya• Praying• Pre-Plenary Session	Ballroom Santika Hotel
08.30 – 10.00	Plenary Session 1 Discussion	Ballroom Santika Hotel
10.00 – 10.30	Coffee Break	Ballroom Santika Hotel
10.30 – 12.00	Plenary Session 2 Discussion	Ballroom Santika Hotel
12.00 – 13.00	Pray & Lunch	floor at Santika Restaurant
13.00 – 16.30	Oral Presentation	Room 1 Room 2 Room 3 Room 4
16.30 – 16.45	Door Prize The best poster and oral presentation appreciation	Ballroom Santika Hotel
16.45 – 17.00	Closing	Ballroom Santika Hotel

X

LIST OF ORAL PRESENTATION

TITLE	PAGE
Public Stigma for Schizophrenic Patient in Kabupaten Malang Lilik Supriati	1
Analysis of Factors Affecting the Ability of Self Management (Self Management) Diabetes Patients Endang Sri P Ningsih, Mahdalena, Syamsul Firdaus.	5
Social Support and Social Network on Self-Concept of Residents Prisons Drugs Case Ii LP Child In Martapura. Syamsul Firdaus, Endang Sri Purwati N, Evy Marlindau	15
Medical Professional's and Patient's Perception on The Importance of Hospice Care Establishment for The Terminal Stage Cancer Patient In Fatmawati Central General Hospital, Jakarta. Okky Rachmad Ngakili, Moch. Prasetyo Mulyanto	21
Activity Level Relationship With Self Esteem in The Elderly Hidayatus Sa'diyah	28
The Overview of The Elderly Lifestyle Profile in Surabaya Christina Y & Spto Dwi Anggoro	36
The Effectiveness Braden Scale and Norton Scale to Pressure Ulcer Risk in Neuro Room Diyah Arini, Imroatul Farida, Intan Ayu	53
The Effect of Endorphin Massage Towards to Decrease Back Pain in Thrid Term of Pregnancy Mother in Dr. Ramelan Navy Hospital Surabaya. Dya Suatrami, Merina W.	59
Use of Concept Model Interpersonal Relationship Stress And Adaptation to Clients After Provision of Social Isolation of Social Skills Training and Family Psychoeducation Use of Concept Model Interpersonal Relationship Stress and Adaptation to Clients After Provision of Social Isolation of Social Skills Training and Family Psychoeducation. Sukma Ayu C.K	65
Health Education Effect on Diabetes Mellitus Type 1 and 2 Compliance Therapy. Dini Mei W, Nur Chabibah	75

TITLE	PAGE
The Effectiveness of Physical Exercise During Hemodialysis Toward Fatigue To Hemodialysis Patients. Dwi Retno Sulistyarningsih, Retno Setyawati, Iwan Ardian	83
Kaili Culture to Reduced Pain and Swelling Because of Filariasis Disease In Puskesmas Baluase. Irsanty Collein, Aminuddin, Andi Fatmawaty Syamsu	91
Internal Factors Affecting Parenting Relationship of Events in Sibling Rivalry Early Childhood Education (Paud) Cemara Ngagel Tirta 31 Surabaya. Lela Nurlela, Dhian Satya R	101
The Effect of Massage on The Sleep Quality Of Elderly. Rian Yuliyana	114
Effect of Music Therapy to Decrease Anxiety Levels in Elderly In Nursing Dormitory One Bhakti Luhur Institution Wisma Tropodo Sidoarjo. Dedi Irawandi	119
The Correlation Between Mother's Stimulation With Child. Mamik R	128
Application of Cooperative Learning Model Type Stad (Student Teams Achievement Divisions) Based on Students Learning Outcomes Jombang. Heni Maryati	134
Effectiveness of Monitor Device PSF 1 To Observation of Breathing Process For Patient Transport in The Ambulance. Abdul Ghofar, Nasrudin	141
Implementation of Activities Daily Living (ADL) Therapy toward Anxiety Before Surgery in Surgical In-patient Ward Dr. Ramelan Hospital Surabaya Dwi Priyantini, Raden Kamaliyatul Adiybah	145
Effect of Dhikr in Recovery Motivation and Blood Glucose of Type 2 Diabetes Mellitus Patients. Rifka Pahlevi	154
Affecting Factors of The Acne Vulgaris Incidence in Bachelor Student of STIKES Hang Tuah Surabaya. Nur Indah Rahmawaty, Qori'ila Saidah, Nisha Dharmayanti R.	162
The Increasing Compliance of Iron Tablet Consumption With Normolity of Hemoglobin Level of Pregnant Women in Second Trimesterat Garum. Fransiska Imavike, Fevriasanty	173
Giving Food Supplement to Recovery With Increasin The Weight in Pre-School Malnutrition. Ririn Probowati	178
Preventing Postpartum Haemorrhage: A Literatur Review. Rini Hayu Lestari	184

TITLE	PAGE
The Effectiveness of ORS And Green Coconut Water. Nur Meity	191
Breastfeeding Mothers With HIV AIDS. Kolifah	200
The Effect of Aseptic Dressing Method on Infusion Phlebitis Prevention Among Hospitalized Patients in Hos pital of Tanjungpinang 2015. Hotmaria Julia Dolok Pasaribu	207
Children Growth Not Influenced By The Quality of Daycare Services. Oswati Hasanah, Sri Romadhona, Riri Novayelinda	219
The Effect of Emergency Level to Nurses Increasing Stress Level at Emergency Room. Ninik Ambar Sari, Candra Perwira	225
The Correction Between Responding Time and Parent Satisfaction Whose Children are Referenced Using Ambulance 118. Dwi Ernawati	232
The Relationship Level of Activity and Transport With Active Participant of Elderly in IHS (Integreted Health Services). Diyah Mutyah	242
Menstrual Disorder Experienced by the Acceptor of Tubectomy in Puskesmas Tambak Rejo Surabaya (Descriptive Observasion). Puji Hastuti, Prasdiana Heny	250
Correction Study of Sustainable Use With IUD KB Anemia on Events IUD Acceptors in BPM Malang Raya 2015. Jenny J Ssondakh, Widya Warastuti	260
Effect of Herbal Therapy on Desreasing Menstrual Pain at Pondok Pesantren Al-Jihad Surabaya. Astrida Budiarti	266

DESCRIPTION OF MENSTRUAL DISORDERS IN ACCEPTORS TUBEKTOMY PUSKESMAS TAMBAKREJO SURABAYA

PujiHastuti¹, Prasdiana Heny Purwanti²
Stikes Hang Tuah Surabaya
Ph_ners79@yahoo.co.id

Abstract

The use of safe contraception such as Method of Operation Women (MOW) today began the public interest because it is a long-term contraception and do not have to perform periodic controls. In Puskesmas Tambak Rejo Surabaya there are free programs from the government to do Method of Operation Women (MOW) so that people, especially women who have children and want to limit the number of children can be admitted to the program. Menstrual disorders are abnormalities that occur during menstruation due to hormonal imbalance, especially women of childbearing age use contraception. This study aimed to describe menstrual disorders experienced by tubal ligation acceptors. Descriptive observational study design. Population acceptor tubektomi in PHC Surabaya Tambak Rejo is 46 used technique *probability sampling* with *simple random sampling* approach as much as 41 respondents. Instrument research using questionnaire. The variables in this study are menstrual disorders. Analysis with descriptive analysis. The results showed menstrual disorders most experienced tubal ligation acceptors are hipomenore as many as 18 people (43.9%), tubal ligation acceptors who experience amenorrhea as many as 16 people (39.0%), and acceptor tubektomi who experience premenstrual syndrome, as many as 15 people (36.6%). The implication of this research is to provide new knowledge to the public about a variety of menstrual disorders. In addition, respondents who complained of menstrual can immediately contact the nearest health facility, so that the nurses can provide appropriate education for prospective acceptor tubal ligation and can health improve and welfare acceptor tubektomi

Keywords: Menstruation, tubal ligation, Women, Hormone

Introduction

Population problems in Indonesia is a large population and uneven distribution, fertility rate and mortality are relatively high in Indonesia. Family planning programs launched by the government of Indonesia to address the fertility rate continues to rise

(Iresnawati, 2012: 280). Family planning is an attempt to measure the number and spacing of children desired by preventing or delaying pregnancy is the contraception (Sulistyawati, 2013: 12). Contraception is divided into 2 of births for a few years and end fertility permanently (Proverawati, et al, 2010 : 1).

Forms of contraception do to end fertility permanently is safe contraception is vasectomy and tubal ligation. Safe contraception in women is known as female tubal ligation or methods of operation (MOW). Research conducted Sri Kustiayati, Hidayat Widjayanegara, Hadyana Sukandarin 2008-2010 about the Female Sexual Function After Tubektomi (Field Study In Surakarta) states that the sexual function of women who have tubal ligation is lower than women without contraception. People in Puskesmas Tambak Rejo many are interested in following tubal ligation because the District Simokerto launched a free program for people who want to do a woman's method of operation. Preliminary studies conducted in Puskesmas Tambak Rejo enough tubal ligation acceptors who complained her menses. Acceptors tubektomi complaining of menstrual most households experience problems ranging from household instability, thinking about school fees, to the desire to have children back so stressful.

Contraceptive use in Indonesia compared to the ASEAN countries in 2005-2012 amounted to 61% (WHO, 2013). For contraceptive users in Indonesia alone in 2012 amounted to 61.9% with the birth rate (*total fertility rate*) of 2.6 % (IDHS, 2012). Use of the method of operation of women (MOW) on a new participant family planning (FP) in Indonesia in 2013 as many as 128 793 people (1.52%) (BKKBN, 2014). Users or acceptors of sterilization of either men or women who are in the largest urban areas amounted to 76.1% for the sterilization of women and 46.6% for male sterilization users (MoH RI, 2013). Preliminary studies conducted on

January 7, 2016 in TambakRejo Surabaya Health Center there are 25 people who do the planning, of the 25 people the 20% who do tubal ligation and complained of irregular menstruation. Results of research conducted EndahSusilowati and EkoPrasetyo in 2012 of Factors Associated With Menstrual Cycle Participants KB Active In DesaJatiKulon, District Jati, Kudus shows that the long use of contraceptives for more than 1 year affect menstrual cycles in acceptors amounted to 97.6%.

Tubal ligation is done with the method of operation to cut and tie the fallopian tubes, causing side effects after surgery. The side effects are bleeding at the site of puncture or tear in the fallopian tubes, bladder injury, until the problem is psychological. Many women who experience regret after tubal ligation to causing psychological problems (Sedesse, 2011). Psychological problems occur because women under going tubal ligation was not satisfied and asked to be restored reproductive function (Hand, 2010). Psychological problems can cause systemic changes in the body, especially the nervous system in the hypothalamus through changes in *endogenous opiates* which may affect expenditures *Gonadotropin Releasing Hormone* (GnRH) (Wirakusumah, et al, 2011). *Gonadotropin Releasing Hormone* (GnRH) secreting FSH and LH (Price & Wilson 2006). Spending *Gonadotropin Releasing Hormone* (GnRH) is not the maximum will affect the secretion of FSH and LH, resulting in the production of estrogen and progesterone disturbed. Impaired hormonal function causing menstrual disorders. Disorders related to

menstruation can cause disturbances in the reproductive process (Kusmiran, 2012). All operative sterilization procedure is intended as a permanent sterilization. Women who use the method of sterilization or tubal ligation surgery is sometimes asked for information about returning sterility. The role of nurses who are at the Center for Public Health in conducting counseling can help acceptor tubektomi and their families to obtain information. Extension granted to acceptor tubal ligation and tubal ligation family may be definition, indications, contraindications, and the impact after a tubal ligation tubal ligation so acceptors can understand the changes resulting from tubal ligation and improve the welfare of the acceptors woman or tubal ligation surgery methods. Based on the above background, it is necessary to study in order to identify the picture on the acceptor menstrual disorders tubal ligation in Tambak Rejo PHC Surabaya.

Method

This research uses descriptive observational design. The population in this study are all acceptor tubektomi in Puskesmas Tambak Rejo Surabaya conducting Operation Methods Women (MOW) for a minimum of 1 year to 46 people. Sampling technique using *probability sampling* with *random sampling* approach *sample* by 41 respondents. The research instrument uses a questionnaire. The variable in this study is menstrual disorders which include disorders of the menstrual cycle, the amount of menstrual blood disorders, and other disorders in menstruation. Data were analyzed using descriptive analysis.

Result

1. Menstrual Cycle Disorders

Table 1.1. Characteristics of respondents by disturbances in the menstrual cycle Tambak Rejo Surabaya Health Center on April 16 - May 16, 2016 (n = 41)

Menstrual Cycle Disorders	Frequency (f)	Percentage (%)
No distractions	18	43.9
Amenorrhoea	16	39.0
oligomenorrhoea	2	4.9
Polimenoire	3	7.3
Metrorrhagia	2	4.9
Total	41	100

2. Total Blood Disorders Menstruation

Table 1.2. Characteristics of respondents by the number of menstrual blood disorders in PHC Tambak Rejo Surabaya on April 16 - May 16, 2016 (n = 41)

Total Blood Disorders Menstruation	Frequency (f)	Percentage (%)
No distractions	11	26.8
Hipomenore	18	43.9
hypermenorrhoea	12	29.3
Total	41	100

3. Other Disorders In Menstruation

Table 1.3. Characteristics of respondents by other menstrual disorders in PHC Tambak Rejo Surabaya on April 16 - May 16, 2016 (n = 41)

Other Disorders In Menstruation	Frequency (f)	Percentage (%)
No distractions	13	31.7
dysmenorrhoea	13	31.7
premenstrual syndrome	15	36.6
Total	41	100

Discussion

Research results in Table 1.1 shows that 41 respondents in TambakRejo PHC Surabaya who did not experience menstrual cycle disorders as many as 18 people (43.9%), respondents who experience amenorrhea as many as 16 people (39.0%), respondents who experienced polimenore as much as 3 people (7.3%), and respondents who have oligomenorrhea and metrorrhagia equal number ie 2 (4.9%). Menstrual cycle disorders experienced by respondents by onset of menstrual disorders after Operation Methods Women (MOW) among other 18 people (43.9%) had no menstrual cycle disorders; 16 people (39.0%) experienced amenorrhea consisting of eight people (53.3%) experienced amenorrhea after MOW for 1 year, 5 (41.7%) experienced amenorrhea after MOW for 2 years, 2 (28.6%) experienced amenorrhea after MOW for 3 years, and 1 person (25.0%) experienced amenorrhea after MOW more than 3 years. Respondents who have oligomenorrhea by 2 people (4.9%) and all disturbed menstrual cycle after Operation Methods Women (MOW) for 3 years.

Menstrual cycle disorders caused can be caused by ovarian arteries clogged and cause hypertension local to the ovaries, resulting in the production of estrogen and progesterone disturbed (Naqvi et al, 2014). Researchers found from the findings that most disturbed menstrual cycle is amenorrhea. In women who use Operation Methods Women (MOW) most experienced amenorrhea. Amenorrhea occurs due to an imbalance of estrogen and progesterone caused by a blockage or cutting the fallopian tube when performing Operation Methods Women (MOW). Estrogen and progesterone produced by the ovaries that

occurs when the ovum maturation will walk past the endometrium into the fallopian tubes. If women use Operation Methods Women (MOW), then the mature ovum will not be able to pass through the fallopian tube toward the endometrium. It will experience an interruption in the menstrual cycle is amenorrhea.

The results of the study in Table 1.2 shows that 41 respondents in Puskesmas Tambak Rejo hipomenore Surabaya who had as many as 18 people (43.9%), respondents who experienced hypermenorrhea as many as 12 people (29.3%), and 11 (26.8%) the rest are not impaired amount of menstrual blood. Respondents were impaired based on the amount of menstrual blood of impaired after Operation Methods Women (MOW) as many as 18 people (43.9%) who experienced such hipomenore 2 (66.7%) experienced hipomenore less than 1 year after MOW, 7 people (46.7%) experienced hipomenore after 1 year do MOW, 4 (33.3%) experienced hipomenore after MOW for 2 years, 2 (28.6) experienced hipomenore for 3 years after the MOW, and 3 (75.0%) experienced hipomenore MOW after more than 3 years. Respondents who experienced hypermenorrhea many as 12 people (29.3%) is 1 person (33.3%) experienced hypermenorrhea less than 1 year after MOW, 3 (20.0%) experienced hypermenorrhea after 1 year do MOW, 4 people (33.3%) experienced hyper menorrhea after 2 years doing MOW, 3 (42.9%) experienced hypermenorrhea after 3 years of doing MOW, and one person (25.0%) experienced hypermenorrhea for more than 3 years after MOW.

Disruption of menstrual blood flow volume can be caused by an imbalance

between the blood supply of the fallopian tubes and ovaries (Ozerkan, et al, 2010). Researchers found that women using the method of operation Women (MOW) will be impaired amount of menstrual blood (hipomenore). This can happen due to GnRH which produce FSH and LH can not go to the ovary to the process of maturation of the ovum. Hormones FSH and LH produced in the hypothalamus will flow to the ovaries along with oxygen and blood flow as a function of blood is carry oxygen throughout the body including the ovaries. If the oxygen carried by the blood to the ovary congested, then the menstrual blood to the acceptor tubektomi experience decline.

Research results in Table 1.3 show that 41 respondents in TambakRejo PHC Surabaya who experience premenstrual syndrome, as many as 15 people (36.6%), respondents who experienced dysmenorrhea people as many as 13 (31.7%), and 13 other people (31, 7%) did not experience other menstrual disorders. Respondents who have other disorders menstrual disorders based on the amount raised after Operation Methods Women (MOW) as many as 15 people (36.6%), is 5 (33.3%) experienced premenstrual syndrome after MOW for 1 year, 7 person (58.3%) experienced premenstrual syndrome after MOW for 2 years, 2 (28.6%) experienced premenstrual syndrome after MOW for 3 years, and 1 person (25.0%) experienced premenstrual syndrome after more than MOW 3 years. Respondents who experience dysmenorrhea as many as 13 people (31.7%) is 1 person (33.3%) experienced dysmenorrhea less than 1 year after MOW, 5 (33.3%) experienced dysmenorrhea after MOW for 1 year, 2

people (16.7%) experienced dysmenorrhea after MOW for 2 years, 2 (28.6%) experienced dysmenorrhea after MOW for 3 years, and 3 (75.0%) experienced dysmenorrhea after MOW more than 3 year.

Researchers argue that if the respondent is not old or new undertaking Operation Methods Women (MOW) can experience menstrual disorders due to the adaptation process of biological and psychological operations fallopian tubes. Biological adaptation in the form of changes in the function of the fallopian tube which was originally to have full functionality to channel ovum from the ovary endometrium is now disrupted because of blockage. Psychological adaptation experienced tubal ligation acceptor can be a problem in itself that happened after Operation Methods Women (MOW) like to think the desire to have children again, problems in the family, to think about problems such as abnormal menstrual previous.

The results are consistent with research conducted Ozerkan, et al (2010) which explains that most women who undergo Operation Methods Women (MOW) has menstrual abnormalities after one year tubal ligation surgery. Naqvi, et al (2014) explains that menstrual changes mostly occurred after tubal ligation for 1 year and menstrual changes can be decreased up to 34% within 3-4 years. The destruction of the fallopian tubes tubal ligation procedure would destroy most mesosalping and change the blood supply to the ovaries. This will reduce the signal *Gonadotropin Releasing Hormone* (GnRH) towards ovarian characterized by a decrease in the number of follicles and corpus luteum function so that the levels of

ovarian hormones are affected and menstrual disorders arise (Tahath, Y and Al-Rayyan, E., 2007).

In terms of age, respondents aged 35-40 years as many as 27 people (65.9%) and respondents aged 41-45 years as many as 14 people (34.1%). Respondents who experience menstrual cycle disorders are mostly experienced by the age of 35-40 years is 12 people (44.4%) experienced amenorrhea, 1 (3.7%) had oligomenorrhea, 2 (7.4%) experienced polimenore, 1 people (3.7%) had metrorrhagia, and 11 (40.7%) had no menstrual cycle disorders. Respondents who experience menstrual cycle disorders aged 41-45 years is 4 people (28.6%) experienced amenorrhea; respondents had oligomenorrhea, polimenore, metrorrhagia each 1 (7.1%); and 7 (50.1%) had no menstrual cycle disorders.

The age of respondents who experience menstrual disorders menstrual blood counts most is 35-40 years as many as 27 people with 11 people (40.7%) experienced hipomenore, 8 (29.6%) experienced hypermenorrhea, and 8 (29.6%) is not impaired amount of menstrual blood. While the number of respondents who experience disruption of menstrual blood at the age of 36-40 for 14 people which 7 (50.0%) experienced hipomenore, 4 (28.6%) experienced hipermeenore, and 3 (21.4%) did not experience the amount of menstrual blood disorders.

Respondents who have other disorders in menstruation is dysmenorrhea which 13 people were 7 (25.9%) experienced dysmenorrhea at the age of 35-40 years and 6 (42.9%) experienced dysmenorrhea at the age of 41-45 years. Respondents who experienced premenstrual syndrome,

as many as 15 people are 9 people (33.3%) experienced premenstrual syndrome at the age of 35-40 years and 6 (42.9%) experienced premenstrual syndrome at the age of 41-45 years. Then as many as 13 people do not experience other menstrual disorders. Respondents who experience dysmenorrhea and premenstrual syndrome, the greatest number are at the age of 35-40 years. Molinc, et al (2010) suggest that the incidence of premenstrual syndrome may be worse at the age above 30 years.

In this study, respondents who often suffer from menstrual disorders aged 35-40 years. Menstrual changes experienced by respondents were also influenced by the age when they make Operation Methods Women (MOW). The younger the woman's mother did Operating Method (MOW), then the risk for the greater menstrual disorders (Shy et al, 1992 in Shobeiri & Atash Khoii, 2005). In addition to menstrual disorders, women who have had Female Operation Methods (MOW) more likely to suffer from dysfunctional uterine bleeding at the age of 35-46 years. Because at that age is a vulnerable age for women and the age most women have dysfunctional uterine bleeding after Operation Methods Women (MOW) or tubal ligation (Zulfiqar & Iftikhar, 2013).

Women older than 35 years and do tubektomi will experience *hot flashes* and menopausal symptoms such as poor sleep quality, night sweats, irritability, and depression. The menopause symptoms worsened in women older age (over 40 years) who have done a tubal ligation (Nichols, et al, 2013). Researchers found the age of the respondent has associated with menstrual disorders. Respondents aged 40-45 years

who have amenorrhea or not menstruation after tubal ligation is feared they are experiencing symptoms of menopause. The older the woman, the greater the risk for premenopausal. In addition bodily functions premenopausal women who are age also decreased, including hormone to its biological function.

Based on respondents who are working, as many as 33 people (80.5%) work as a housewife, 6 (14.6%) working as a private, 1 (2.4%) worked as an entrepreneur, and one person (2, 4%) more are working as temporary teacher. Respondents who experience menstrual cycle disorders based on the work that 16 people have amenorrhea with details of 13 people (39.4%) work as a housewife, 2 (33.3%) work as a private, and 1 (100.0%) working as self-employed. Respondents who have oligomenorrhea as much as 2 ie 1 (3.0%) worked as households and 1 (16.7%) working as a private. Respondents who experienced polimenore as many as 3 people (9.1%) who works as a housewife. Respondents who experienced metrorrhagia much as 2 ie 1 (3.0%) worked as a housewife and 1 (100.0%) more are working as a temporary teacher. then 18 people experience no disruption of the menstrual cycle.

Respondents who experience disruption of menstrual blood volume based on the work that 18 people have hipomenore with details 13 people (39.4%) work as a housewife, and 5 (83.3%) working as a private. Respondents who experienced hypermenorrhea many as 12 people with 11 people (33.3%) worked as a housewife and 1 (100.0%) working as a private. A further 11 people are not impaired amount of menstrual blood. Works of respondents who experienced

other disturbances in the menstrual mostly experienced by mothers house as many as 33 people which is 10 people (30.33%) experienced dysmenorrhea, 14 people (42.4%) experienced premenstrual syndrome, and 9 (27.3%) did not other menstrual disorders.

Respondents who worked as private employees as much as six persons, namely 3 (50.0%) experienced dysmenorrhea and 3 (50.0%) did not experience other menstrual disorders. Respondents who worked as a temporary teacher as much as 1 (100.0%) experiencing premenstrual syndrome, and 1 (100.0%) worked as an entrepreneur who did not experience other menstrual disorders. Physical activity of moderate to severe may inhibit *gonadotropin-releasing hormone* (GnRH) which can decrease serum ekstrogen (Hoyer, et al, 2013). Researchers found respondents who worked as a housewife has to take care of all household purposes eg cooking, washing, to educate their children will experience fatigue mereka.sehingga its own that can trigger systemic changes in the body including hormonal changes of menstruation.

Operation method Women (MOW) or tubal ligation is the most effective contraceptive method for use of a permanent and highly effective (Edelman, et al, 2013). Tubektomi also raised concerns about the potential disruption after tubal ligation both biologically and psychologically including regret after tubal ligation procedure because the procedure is generally *irreversible*. Abnormalities of menstruation is a disorder that is often experienced by women who have tubal ligation. Menstrual disorders caused after a tubal ligation due to blood circulation

in and around the fallopian tubes and ovaries, the pressure on the nerve, and adhesion intrapelvic noncurrent (Satoh, 1993 in Palli, 2013). Tahath, Yousef and Al-Rayyan (2007) analyzed hormonal, endometrial biopsy, and observe the changes of menstruation. The result they found a decrease midluteal phase estradiol (E2), progesterone (P), and *luteinizing hormone* (LH) in women who do a tubal ligation.

Different methods used or the type of surgery can destroy other parts of the fallopian tube and the blood flow (Naqvi et al, 2014). Tubal ligation procedure can be done by Irwing, Pomeroy, Uchida, Parkland method, and using catheterization unipolar, bipolar up using various clips. Wilcox, et al (1992 in Ozerkan, et al, 2010) observed 5070 patients over 5 years and found that the method per clip lead to menstrual pain and bleeding are more in 33% of patients. Additionally longest menstrual period was found in the case of unipolar electrocoagulation, and the shortest menstrual period experienced by women who use per clip. Parsanezhad, et al (2008) suggested that the technique Pomeroy, rings, and methods of unipolar cause more damage to the utero-tubal and utero-ovarian. The level of damage to blood vessels peritubal and ovaries can cause menstrual abnormalities after tubal ligation.

Respondents who experience amenorrhea as many as five people consisting of 4 people have amenorrhea and still give milk, and 1 person amenorrhea after Operation Methods Women (MOW). While 35 of the respondents had more than one menstrual disorders. Women who do Operation Methods Women (MOW) or tubal ligation generally do not impact on

breastfeeding her baby because tubal ligation is a safe method of contraception (Berens, et al, 2015). The cause of amenorrhea in tubal ligation acceptors who are still breastfeeding may be affected by current drugs do Surgery Methods Women (MOW). Women who choose to do tubal ligation shortly after birth the baby will be given general anesthesia by medical teams and activities will continue to breastfeed after the operation performed by the medical team record should make the best decisions on the selection of safe anesthesia. The medical team should avoid the use of meperidine (Demerol) or the use of diazepam (Valium) long-term anesthesia when performing tubal ligation. However, local anesthetics, benzodiazepines (eg Ativan), muscle relaxants, inhalation agents can be used in the procedure of tubal ligation surgery and anesthesia is quite safe (Edelman, et al, 2013).

Conclusion

The results of the research that has been done in Puskesmas Tambak Rejo Surabaya on April 16 - May 16, 2016 can be drawn conclusions menstrual disorders experienced by the acceptor tubektomi largely composed of several disorders that menstrual cycle disturbances, impaired amount of menstrual blood, and other disorders in menstruation. Menstrual cycle disorders experienced by the majority of acceptors tubektomi namely amenorrhea, impaired amount of menstrual blood is experienced mostly acceptor tubektomi namely hipomenore, while other disturbances in the menstrual experienced mostly acceptor tubektomi namely premenstrual syndrome, menstrual

disorders is because the blood circulation in and around Fallopian tubes and ovaries is not smooth. Smooth blood circulation is caused by the destruction of most of mesosalping the fallopian tube when tubal ligation procedure. This matter will cause signal *Releasing Gonadotropin Hormone (GnRH)* to the ovaries decreases were characterized by a decrease in the number of follicles and corpus luteum function so that menstruation becomes impaired.

References

- Berrens, P., dkk. (2013). *Contraception During Breastfeeding*. <http://online.liebertpub.com/doi/pdf/10.1089/bfm.2015.9999sitasitanggal> 15 Juni 2016 pukul 03.10 WIB
- Edelman, A., dkk. (2013). *Benefits And Risks of Sterilization*. <http://pdfs.journal.lww.com>. sitasi tanggal 9 Juni 2016 pukul 21.00 WIB
- Handayani, S. (2010). *Buku Ajar Pelayanan Keluarga Berencana*. Yogyakarta: PustakaRihama
- Hoyer, J., dkk. (2013). *Menstrual Cycle Phase Modulates Emotional Conflict Processing In Women With And Without Premenstrual Syndrome (PMS) - A Pilot Study*. [http://journals.plos.org/pone.0059780sitasitanggal](http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0059780sitasitanggal) 10 Juni 2016 pukul 03.45 WIB
- Kusmiran, E. (2012). *Kesehatan Reproduksi Remaja dan Wanita*. Jakarta: Salemba Medika
- Kustiyah, S., Widjayanegara, H., Sukandar, H. (2015). *Fungsi Seksual Wanita Pasca Tubektomi (Studi Lapangan di Kota Surakarta)*. *GAS'ER Vol. XII No. 1 Februari 2015*
- Moline, Margaret, L., dkk. (2010). *Premenstrual Dysphoric Disorder: A Guide for Patients and Families*, https://womensmentalhealth.org/wp-content/uploads/2008/04/pmdd_guide.pdfsitasitanggal 30 Mei 2016 pukul 16.15 WIB
- Naqvi, Syeda., dkk. (2014) *Spectrum Of Menstrual Problems After Tubal Ligation*, http://pafmj.org/pdfs/june-2014/Article_35.pdfsitasitanggal 30 Mei 2016 pukul 15.50 WIB
- Nichols, H, B., dkk. (2013). *Tubal Ligation In Relation To Menopausal Symptoms And Breast Cancer Risk*. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3778289/pdf/bjc2013433a.pdf>sitasitanggal 9 Juni 2016 pukul 08.00 WIB
- Ozerkan, Kemal., dkk. (2010). *Menstrual Pattern Following Tubal Sterilization*, <http://www.medscimonit.com/download/index/idArt/878502sitasitanggal> 30 Mei 2016 pukul 15.00 WIB
- Palli, S, B dan Akkupalli, V, L. (2013). *A Study To Evaluate The Relationship Between Socio-Economic, Demographic Characteristics And Complication Of Acceptors In Double Puncture Laparoscopic And Conventional Tubal Ligation Procedures*. <http://www.scopemed.org/fulltextpdf.php?mno=32107sitasitanggal> 9 Juni 2016 pukul 13.20 WIB
- Parsanezhad, ME., dkk. (2008). *Menstrual Abnormalities And Pain After Five Tubal Sterilization Methods: A Randomized Controlled Trial*. <http://ijms.sums.ac.ir/index.php/IJMS/article/view/2432/809sitasitanggal> 9 Juni 2016 pukul 08.20 WIB
- Proverawati, A., Islaely, A., Aspuah, S. (2010). *Panduan Memilih Kontrasepsi*. Yogyakarta: NuhaMedika
- Sedesse, P. (2011). *Long Term Side Effect of Having Tubes Tied and The Effect on Hormone Levels*, <http://www.explain-health.com/tubal-ligationsitasitanggal> 2 Februari 2016 pukul 19.20 WIB
- Shobairi, M, J &AtashKhoii, S. (2005). *The Risk Of Menstrual Abnormalities After Tubal Sterilization: A Case Control Study*, <http://bmcwomenshealth.biomedcentral.com/>

- articles/10.1186/1472-6874-5-5sitasitanggal 8 Juni 2016 pukul 19.00 WIB
- Susilowati, E., Prasetro, E. (2015). Faktor-faktor yang Berhubungan dengan Siklus Menstruasi Peserta KB Aktif di Desa Jati Kulon Kecamatan Jati Kabupaten Kudus. *JIKK Vol. 6 No. 1 Januari 2015 Page 79-96*
- Tahat, Y, A & Al-Rayyan, E, S. (2007). *What Happens After Tubal Sterilization: A Jordanian Experience*. <http://rmsjournal.org/Articles/635779038627642238.pdf>sitasitanggal 8 Juni 2016 pukul 19.15 WIB
- Wirakusumah, E., Mose, C., Handono, B. (2011). *Obstetri Fisiologi: Ilmu Kesehatan Reproduksi Edisi: 2*. Jakarta: EGC
- Zulfiqar, B dan Iftikhar, R. (2013). *Late Gynaecological Complications Following Pomeroy Method Of Mini Laparotomy For Tubal Ligation*. <http://www.jsp.org.pk/Issues/JSP%2018%20%284%29%20Oct-Dec%202013%20PDF/Bushra%20Zulfiqar%20OA.pdf>sitasitanggal 9 Juni 2016 pukul 16.50 WIB



Certificate



Puji Hastuti, M.Kep.,Ns

for his/her contribution as

Oral Presenter

THE 2nd INTERNATIONAL NURSING CONFERENCE (INC) 2016
"Developing Cross-Cultural Understanding and Behavior in Nursing Care Services"

In Surabaya, Indonesia November 5th, 2016

Accredited based on decree of Indonesia National Nurses Association

No: 1036/ DPP.PPNI / SK / K.S / XI / 2016

Speaker 3 SKP, Participant 3 SKP, Moderator 2 SKP, Committee 2 SKP.



Wiwie K. Liestyningrum, S.Kp., M.Kep.
The Head of STIKES Hang Tuah Surabaya



Puji Hastuti, S.Kep., Ns., M.Kep.
Chairperson of Organizing Committee